

UNIVERSITY OF COPENHAGEN



## Comparative Genomics of *Vibrio cholerae* O1 Isolated from Cholera Patients in Bangladesh

Hossain, Zenat Zebin; Leekitcharoenphon, Pimlapas; Dalsgaard, Anders; Sultana, Rokaia; Begum, Anowara ; Jensen, Peter Kjær Mackie; Hendriksen, Rene S.

*Publication date:*  
2017

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Hossain, Z. Z., Leekitcharoenphon, P., Dalsgaard, A., Sultana, R., Begum, A., Jensen, P. K. M., & Hendriksen, R. S. (2017). *Comparative Genomics of Vibrio cholerae* O1 Isolated from Cholera Patients in Bangladesh. Poster session presented at International Conference On Genomics Nanotech & Bio engineering , Dhaka, Bangladesh.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/314564820>

## Comparative Genomics of *Vibrio cholerae* O1 Isolated from Cholera Patients in Bangladesh

Conference Paper · May 2017



CITATIONS  
0

7 authors, including:




**Zenat Zebin Hossain**  
University of Copenhagen  
10 PUBLICATIONS 0 CITATIONS  
[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:

-  Excreta reuse in Vietnam [View project](#)
-  The importance of water quality on the risk of diarrhea: A prospective cohort analysis of people living in Southern Punjab, Pakistan [View project](#)

READS  
91



**Pimlapas Leekitcharoenphon**  
Technical University of Denmark  
76 PUBLICATIONS 700 CITATIONS  
[SEE PROFILE](#)





# Comparative Genomics of *Vibrio cholerae* O1 Isolated from Cholera Patients in Bangladesh

Zenat Zebin Hossain<sup>1,2</sup>, Pimlapas Leekitcharoenphon<sup>3</sup>, Anders Dalsgaard<sup>4</sup>, Rokaia Sultana<sup>1</sup>, Anowara Begum<sup>1</sup>, Peter Kjær Mackie Jensen<sup>2,5</sup>, Rene S. Hendriksen<sup>3</sup>

<sup>1</sup>Department of Microbiology, University of Dhaka, Dhaka 1000, Bangladesh; <sup>2</sup>Institute of Public Health, University of Copenhagen, Copenhagen 1014, Denmark; <sup>3</sup>National Food Institute, Technical University of Denmark, Research Group for Genomic Epidemiology, WHO Collaborating Center for Antimicrobial Resistance in Food borne Pathogens and Genomics and European Union Reference Laboratory for Antimicrobial Resistance, Kgs. Lyngby, Denmark; <sup>4</sup>Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, 1870 Frederiksberg C, Copenhagen, Denmark; <sup>5</sup>Copenhagen Centre for Disaster Research, University of Copenhagen, Copenhagen 1014, Denmark

## CONTACT

Contact information  
email: zenat@sund.ku.dk; zenatzebinorin@yahoo.com  
Department of Microbiology, University of Dhaka & Department of Public Health, University of Copenhagen  
Ph: +8801785636044

## BACKGROUND

- Cholera remains an endemic disease in Bangladesh where this disease occurs in seasonal regularity with more than 100,000 cases annually (Ali et al; 2015). Recently, the severity of the disease has significantly increased in urban area since the emergence of the new variant of *Vibrio cholerae* O1 El Tor (Chowdhury et al; 2015).
- Molecular changes in *V. cholerae* have to be closely monitored as this information may help in understanding the changing genetic features of this pathogen in relation to the epidemiology of cholera.
- In this study, Whole Genome Sequencing (WGS) was utilized to investigate the current genomic profile of *V. cholerae* O1 strains, isolated from symptomatic patients in the low-income urban area of Arichpur, Dhaka, Bangladesh.

## METHODS AND MATERIALS

During October 2015, three *V. cholerae* O1 strains (VC-1, 2 and 3) were isolated from rectal swabs of two patients living in households 588 m apart. One of the two patients was co-infected with two *V. cholerae* strains (VC-1 and VC-3). Major virulence factors, biotype and antimicrobial resistance genes were identified by WGS. A global phylogenetic tree was inferred using genome wide SNPs (Single Nucleotide Polymorphism) analysis

### Sequencing Platform and Online Bioinformatics Tools




Table 1: Strain information

V. cholerae strain ID	Source	City/Country	Accession No.	Serogroup	Serotype	Biotype	Species-Serogroup	Biotype-specific			
								ompW	rfbV-O1	wbfZ-O139	ctxB
VC_1	Rectal swab	Dhaka/Bangladesh	ERS1484954	O1	Inaba	El Tor variant	+	+	-	CC*	ET*
VC_2	Rectal swab	Dhaka/Bangladesh	ERS1484955	O1	Inaba	El Tor variant	+	+	-	CC	ET
VC_3	Rectal swab	Dhaka/Bangladesh	ERS1484956	O1	Inaba	El Tor variant	+	+	-	CC	ET

Table 2: Virulence profile

VC strain ID	Virulence genes											Gene for 7th pandemic	intSXT gene (mobile genetic element)	Class I integron (intl)
	ctxA	ctxB	tcpA	zot	ace	hlyA	stn	chxA	rtxA	toxR	ompU	mshA	TTSS	
VC_1	+	+	+	+	+	+	-	-	+	+	+	+	-	+
VC_2	+	+	+	+	+	+	-	-	+	+	+	+	-	+
VC_3	+	+	+	+	-	+	-	-	+	+	+	+	-	+

Table 3: Resistance gene profile

Isolate ID	Antimicrobial Agent	Resistance genes
VC-1	Aminoglycoside	strB
	Phenicol	catB9
	Trimethoprim	dfrA1
VC-2	Aminoglycoside	strA, strB
	Phenicol	catB9
	Sulphonamide	sul2
VC-3	Trimethoprim	dfrA1
	Aminoglycoside	strA, strB
	Phenicol	catB9

## RESULTS

- All three *V. cholerae* strains were El Tor variant of *Vibrio cholerae* O1 carrying standard classical ctxB genotype.
- The three strains are strictly clonal, 2 SNPs difference of VC-1 with VC-2 and 3, no SNP difference between VC-2 and 3. The rational behind the low number of SNPs in these Vc strains might be that the strains were isolated within a limited time frame (11 days apart) and from local outbreaks (two households are in the same community).
- The closest neighboring genomes in SNP based global phylogenetic tree were differed in 11 and 12 SNPs which were epidemic clones of *V. cholerae* O1 isolated in 2010 from Pakistan and Bangladesh respectively.
- All strains harbored multidrug resistance genes for aminoglycosides, phenicol, sulphonamide, and trimethoprim except VC-1 that lacked sulphonamide resistance genes. They lacked the integrase gene of the class I integron (intl) but harboured the SXT element (intSXT).
- The in silico MultiLocus Sequence Typing (MLST) revealed that these strains belonged to sequence type 69 which is similar to the pandemic strains (N16961 and MO45) and the Haitian epidemic strain (2010EL1786).




Figure 1: SNP matrix of the three study strains




Figure 2: Global phylogenetic tree constructed with 352 database strains based on SNPs (Regions highlighted in colors)

## CONCLUSIONS

- In this study, we used WGS and bioinformatics tools to analyze current genetic traits of clinical *V. cholerae* O1 circulating in urban household clusters of Bangladesh.
- The study contributed to understanding the epidemiology of *V. cholerae* in household patients in Bangladesh that ultimately can be applied for control measures and management of the emerging pandemic cholera in Bangladesh.

## REFERENCES

- Ali, M., Nelson, A. R., Lopez, A. L., Sack, D. A., 2015. Updated global burden of cholera in endemic countries. PLoS Negl Trop Dis 9, e0003832.
- Chowdhury F., Kuchta A., Khan A.I. et al; 2015. The increased severity in patients presenting to hospital with diarrhea in Dhaka, Bangladesh since the emergence of the hybrid strain of Vibrio cholerae O1 is not unique to cholera patients . Int J Infect Dis.

## ACKNOWLEDGEMENT

This study was funded by the project entitled “Combating Cholera Caused by Climate Change in Bangladesh, CS” (Grant no. 12-040KU) from Danish International Development Agency.